

Specificity and randomness in the visual cortex

Kenichi Ohki and R Clay Reid

Research on the functional anatomy of visual cortical circuits has recently zoomed in from the macroscopic level to the microscopic. High-resolution functional imaging has revealed that the functional architecture of orientation maps in higher mammals is built with single-cell precision. By contrast, orientation selectivity in rodents is dispersed on visual cortex in a salt-and-pepper fashion, despite highly tuned visual responses. Recent studies of synaptic physiology indicate that there are disjoint subnetworks of interconnected cells in the rodent visual cortex. These intermingled subnetworks, described *in vitro*, may relate to the intermingled ensembles of cells tuned to different orientations, described *in vivo*. This hypothesis may soon be tested with new anatomic techniques that promise to reveal the detailed wiring diagram of cortical circuits.

Addresses

Department of Neurobiology, Harvard Medical School,
220 Longwood Avenue, Boston, MA 02115, USA

Corresponding author: Reid, R Clay (clay_reid@hms.harvard.edu)

Current Opinion in Neurobiology 2007, **17**:401–407

This review comes from a themed issue on
Sensory systems
Edited by Peter Mombaerts and Tony Zador

Available online 27th August 2007

0959-4388/\$ – see front matter

© 2007 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.conb.2007.07.007](https://doi.org/10.1016/j.conb.2007.07.007)

Introduction

Over the past 50 years, the visual cortex has served as a model system for the study of cerebral cortical circuits. Several themes have dominated this literature: hierarchical versus recurrent processing, specific versus random synaptic connectivity, and functional architecture versus intermingling of response types. A classical view of visual cortical processing [1] has concentrated on one side of each of these dichotomies that the visual cortex is best understood as a hierarchical system, whose receptive fields are created through specific connections, within a framework of crystalline functional architecture. At various times, however, each of these views has been called into question.

Orientation selectivity and hierarchy

The hierarchical model of visual processing was proposed at the beginning of modern studies of the visual cortex, when it was proposed that selectivity for stimulus

orientation emerges from the specific connections from thalamus to cortical simple cells [2]. In the cat visual cortex, there is evidence that multiple thalamic afferents, each of which is substantially unoriented, add together to produce a strongly oriented afferent input to simple cells in layer 4 [3,4]. This evidence comes from cross-correlation studies [5] as well as studies of orientation selectivity of thalamo-recipient cells when the intracortical circuit has been silenced [6–8].

While an increasing body of evidence has shown that thalamo-cortical connections are related to the establishment of orientation selectivity, the role of intracortical connections has been much more difficult to study. Cross-correlation techniques have shown that simple cells in layer 4 connect to iso-orientation complex cells [9], but the detailed logic of these connections is not clear. There is new evidence that simple cells predominate in layer 4 and complex cells in other cortical layers [10], but the relative importance of feedforward, recurrent, and feedback connections in the cortical circuit is still vigorously debated. Without newer tools to study the relationship between intracortical synaptic connections and visual response properties, it will be, however, difficult to resolve these issues.

While there is strong physiological evidence for the role of thalamo-cortical input in orientation selectivity, a more direct anatomical proof has been elusive. In some species, however, the main thalamo-recipient cells in layer 4 are not orientation selective, but cells in the next stage—layer 2/3 are orientation selective. Their orientation selectivity must develop through intracortical connections from layer 4 to layer 2/3. In an elegant recent study of one such species, the tree shrew, an elongated topographic organization of connections from layer 4 to layer 2/3 correlated well with the emergence of orientation specificity [11].

Coarse-grained and fine-grained specificity of intracortical connections

Since the Hubel and Wiesel's model was first put forth, the specificity of the anatomical connections has been studied at increasingly fine levels of detail: from areal maps to functional or laminar architecture, fine-scale geometry, and finally individual synaptic connections. At the coarsest levels are cortical regions, which often have large-scale maps, such as the retinotopy of the visual cortex. At the next level is functional architecture, in which different features can be segregated at a scale of tens to hundreds of microns [1]. The afferent connections to cortex often respect these functional boundaries, such

as in the ocular dominance columns in the input layers of primate visual cortex [12], or in the cortico-cortical projections between different functional compartments [13,14]. Many studies have also demonstrated specificity in intracortical connections, either between cortical layers [15,16] or in long-range connections within layers [17,18].

The next finer scale has been termed the *geometrical* level at which the close proximity of axons and dendrites are considered on the scale of $\sim 0.5\text{--}2\ \mu\text{m}$ [19,20,21]. At this level and the next – that of actual synaptic connections – the debate about specificity has been couched in terms of *Peters' rule* [19,22,23], that is, axons make connections randomly in direct proportion to the occurrence of all synaptic targets in the adjacent neuropil, with no local specificity.


Recently, several fine-scale geometrical analyses have been performed for cortical circuits. For pyramidal neurons in somatosensory cortex, at least one direct apposition of axons and dendrites (regardless of the existence of a synapse) was observed for every pair of pyramidal neurons sharing the same cortical column within $300\ \mu\text{m}$ [20,21]. This supports Peters' rule for the geometrical contact between axons and dendrites. But this level can only reveal potential connectivity: Actual synaptic boutons were found in only a fraction of these potential connectivities. The number of synaptic boutons was correlated to the synaptic responses of pairs of neurons [20].

A version of Peters' rule was tested directly by comparing geometrical and functional connectivities [24] in barrel cortex. Functional connectivity could not be predicted solely from the overlap of dendrites and axonal arbors, but much of the variations could be explained when other factors were considered. The probability of finding synaptic connections depended on the class and laminar location of each neuron. Furthermore, synaptic connectivity also depended on a neuron's location in the cortical map, that is, whether it was in a barrel or a septum.

Physiological studies of synaptic connectivity have also shown that neurons are not connected randomly to potential targets. Dual intracellular recording studies in slices have shown that neurons are synaptically connected to only a small fraction of their neighbors, and that such synaptic connections are specific to cell types [25–29]. Even within a single-cell type, pyramidal cells, a recent statistical analysis of simultaneous recordings has shown that connections are far more clustered than expected from a random network [30].

Two recent studies of rodent visual cortex have revealed the existence of spatially overlapping but distinct subnetworks of cells. In one study of layer 2/3 pyramidal neurons [31], lateral connections between cells were frequently observed when they received common inputs from layer 4 (Figure 1c). When they did not receive common inputs, even adjacent neurons were rarely connected to each other. The existence of subnetworks was reinforced with a subsequent study of two classes of layer 2/3 inhibitory

Figure 1



Single-cell resolution orientation maps from (a) a pinwheel in cat visual cortex [42] and (b) rat visual cortex [38] obtained with *in vivo* two-photon calcium imaging. One side is $300\ \mu\text{m}$. Cells are colored according to their preferred orientation. In (a), ~ 1000 cells from nine different depths are overlaid. Cells are arranged up to the very center of the pinwheel in cat visual cortex. In (b), cells in one depth of rat visual cortex are displayed. Even neighboring cells are tuned to different orientations. (c) In rat visual cortex, relatively independent subnetworks are embedded in larger scale functional architecture [31]. Excitatory connections from layer 4 to layer 2/3 and within layer 2/3 define subnetworks of selectively interconnected neurons (red or blue). The excitation from layer 5 (gray triangles) and inhibition from layers 2/3 and 4 adaptive interneurons (IN, gray ovals) do not respect the subnetworks.

interneurons: fast-spiking and adapting [32]. Pairs of pyramidal cells and fast-spiking (FS) interneurons are likely to be reciprocally connected only when they share common input. Fast-spiking cells are thus specifically connected with pyramidal cell subnetworks. By contrast, connections from adapting interneurons were not specific. The functional role of these cortical subnetworks is not immediately apparent, but new imaging techniques hold the promise of relating network connectivity and function.

Cellular imaging *in vivo*: the fine-scale architecture of visual processing

Early extracellular recordings of single cells in visual cortex have shown that neighboring neurons often exhibit similar response properties such as ocular dominance and orientation selectivity. Functional groups are arranged in columns, as demonstrated by penetrations made normal to the cortical surface. Tangential penetrations, on the contrary, show regular variations and predictable shifts, suggestive of systematic maps [2]. The orderly progression of receptive-field types across the cortical surface along with the similarity within columns is known as the functional architecture of visual cortex. Optical imaging revolutionized the study of functional architecture by showing the overall geometry of functional maps [33,34]. The spatial resolution of conventional optical imaging (>100 μm) is well matched to most features of cortical maps, but it is inadequate to examine the fine-scale features, such as orientation pinwheels and direction fractures [35].

In vivo two-photon calcium imaging of population of neurons is a new technique [36] for single-cell resolution functional imaging. *In vivo* calcium imaging in the cerebral cortex had only been achieved by intracellular labeling of single neurons [37], until a method for simultaneously loading many neurons with a cell-permeant (AM-ester) form of calcium indicators was developed recently [36]. Using this method, it has become possible to map the response selectivity of all neurons in a local cortical volume (300–600 μm in diameter) at single-cell resolution [38**]. This technique is unique in its ability to determine not only the physiological response of hundreds of cells simultaneously but also their precise locations in the cortical circuit.

Sharp functional borders

Sharp transitions across functional borders have been observed from the outset of the discovery of functional columns by Powell and Mountcastle [39]. More extensive studies revealed honeycomb-like structures of discrete functional columns in the somatosensory cortex [40]. In the visual cortex, preferred orientation changes smoothly, except for occasional abrupt discontinuities in the sequence [41].

Two-photon calcium imaging has recently demonstrated the extraordinary precision of the functional architecture

of orientation and direction selectivity in cat visual cortex [38**,42]. In area 18 of cat visual cortex, iso-orientation domains were known to contain subregions with opposite preferred directions [43]. Two-photon calcium imaging demonstrated that neurons with opposite preferences for stimulus direction were segregated by remarkably precise 'direction fractures', with columnar borders one to two cells wide [38**].

Another class of singularities in the visual cortical map is the orientation pinwheel center [33,34]. Conventional optical imaging first revealed these pinwheels, but the technique lacked the spatial resolution to determine the response properties and arrangement of cells near pinwheel centers. Electrophysiological recordings later demonstrated sharply selective neurons near pinwheel centers [44,45], but it remained unclear whether they were arranged randomly or in an orderly fashion. Two-photon calcium imaging revealed that pinwheel centers are highly ordered: Neurons selective to different orientations are clearly segregated even in the very center of pinwheels [42] (Figure 1a). Thus, pinwheel centers truly represent singularities in the visual cortical map.

The finding of sharp discontinuities in cortical maps raises more questions than answers. The discontinuities are more precise than the spatial scale of the dendritic trees of cortical neurons, so one can ask: What are the mechanisms underlying this precision? Several possibilities are (1) that selective connections between individual neurons with similar tuning dictate the receptive-field properties on either side of the border, (2) that smaller anatomical features, such as bundles of apical dendrites [46,47] or inputs to proximal dendrites, are functionally important, or (3) nonlinear input–output transformation, such as the threshold for spike generation, could sharpen the tuning of neurons around functional discontinuities [45,48]. Single-cell calcium imaging combined with other techniques might distinguish these different possibilities. For example, the relation between anatomy and physiology could be obtained by labeling cells to reveal their transmitters or other molecular markers, their projection patterns, or their detailed dendritic and axonal morphologies.

How do functional features such as directional fractures develop? Orientation maps exist at the time of eye opening [49], but direction maps appear several days after eye opening and do not develop in animals deprived of visual experience [50]. The emergence of the directional map is experience dependent and selective to experience. In a recent study, visually naïve ferrets were exposed to bars of a single orientation moving in two opposite directions, and direction selective patches were induced very rapidly, after 12–18 h of visual experience (Li *et al.*, abstract in Soc Neurosci Abstr 2006, 619.6). It would be very interesting to observe the rapid plasticity of direction fractures at a single-cell level.

Response selectivity without functional maps

We saw that orientation maps in higher mammals are crystalline and organized at the level of single cell, but what is the significance of these maps? Hubel and Wiesel [51] proposed that

It seems reasonable to suppose that the closer cells are in a nervous structure the better their chances will be of having interconnexions or of sharing connexions: there is at least a certain economy in having cells that share connexions close to one another.

But the functional architecture of the cortex is not always necessary to obtain highly selective responses [38^{••},52[•],53] (reviewed in reference [54[•]]). In visual cortex of mice [55–57] and rats [58–60], no evidence has been found for orientation maps, though neurons are sharply tuned to orientation. Two-photon calcium imaging confirmed that, in the rat, a mixed salt-and-pepper organization can still yield strong stimulus selectivity [38^{••}] (Figure 1b). A similar salt-and-pepper organization was found in mouse visual cortex [61,62[•]] (Mrsic-Flogel *et al.*, abstract in Soc Neurosci Abstr 2005, 742.6).

The mixed, salt-and-pepper arrangement of preferred orientation in rodents [38^{••}] argues for specific connectivity between neurons. The orientation tuning of any given neuron cannot be achieved by simply ‘going along for the ride’ within a neighborhood of similarly tuned cells. Strong orientation tuning in rodents instead is probably the result of specific mechanisms such as selective connections among cells with similar response properties, which may form subnetworks of neurons [31[•]].

But what explains the lack of orientation maps in rodents? Rodents could have too small of a visual cortex to have an organized map. Or they could have too low of visual acuity to necessitate one. However, the gray squirrel, a rodent with a relatively large visual cortex (larger than tree shrew; comparable with ferret) and good visual acuity (comparable with tree shrew), does not have orientation maps [52[•],63].

This may be explained by differences in intracortical circuits and wiring length minimization [64]. As might be expected from the above quotation from Hubel and Wiesel [51], the existence of functional architecture in cat visual cortex may be explained by wiring length minimization. But recent theoretical studies have suggested that the *absence* of functional architecture in rodent visual cortex could be more efficient for a specific class of functional wiring diagrams of a cortical circuit [64]. We currently do not know how differ the wiring diagrams of rodent and cat visual cortex. Such local connectivity can be studied only when microscopic functional imaging is combined with higher resolution anatomical techniques that elucidate neuronal morphology and even individual synaptic connections.

New techniques for mapping connections in cortical microcircuits

While the new results in two-photon functional imaging raise many questions, many of them might be addressed by new anatomical and physiological techniques for analyzing neural circuits. At one extreme, automated methods in serial-section electron microscopy [65] have renewed the hope of making large-scale wiring diagrams of cortical circuits. In the shorter term, the ability to trace subnetworks of neurons can be traced by means of viruses [66–68]. Most notably, a newly modified rabies virus has been demonstrated to label specifically all of the neurons that provide monosynaptic input to a single-target neuron [69^{••}]. Combining single-cell resolution functional mapping with these anatomical techniques will give increasingly complete functional and anatomical pictures of the cortical circuit.

The evolution of genetic tools in mice has fostered an unprecedented opportunity to study neurons in living animals. The ability to stably label specific subsets of neurons with genetically-encoded reporters allows the dissection of cell-type-specific function in the dauntingly complex cortical circuit. Several groups have generated mouse lines with subsets of neurons labeled with fluorescent proteins using various promoters [70–74]. By combining power of two-photon calcium imaging and the cell-type-specific labeling afforded by mouse genetics, the complexity of the cortical network can be studied by examining cell-type-specific function in intact cortical circuits [65]. This approach can lead to a more complete understanding of sensory physiology of interneurons and their role in cortical information processing.

Finally, there is an increasingly rich arsenal of techniques for influencing the activity of cortical circuit with cellular resolution [75,76,77[•]]. Cell-type-specific suppression [78] may reveal the function of each cell type in cortical processing. Optical control of neural activity with genetically introduced light-gated channels, such as channelrhodopsin-2 and halorhodopsin [79^{••},80[•],81[•]], is opening up a way to manipulate neural activity at millisecond resolution. These techniques in combination with calcium imaging should soon allow us to turn on and off individual neurons while observing circuit activity in response to sensory stimulation.

Conclusion

There have been considerable debates on the specificity and randomness of the connections in the cerebral cortex. Recently, cellular-resolution imaging of cortical circuits *in vivo* has started to reveal the functional micro-organization of visual cortex. Results in rodent visual cortex, which displays highly selective visual responses without functional architecture, are consistent with specific connections among subnetworks of cells. Conversely, an increasing number of *in vitro* studies have revealed

specific connections and subnetworks but do not reveal their function. Over the coming years, the relationship between functional micro-organization and fine-scale specific connections should be revealed by combination of functional imaging and new anatomical techniques.

Acknowledgements

This work is supported by NIH grant RO1 EY10115. We thank Vincent Bonin and Mark Histed for comments on the manuscript.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Hubel DH, Wiesel TN: **Ferrier lecture. Functional architecture of macaque monkey visual cortex.** *Proc R Soc Lond B Biol Sci* 1977, **198**:1-59.
 2. Hubel DH, Wiesel TN: **Receptive fields, binocular interaction and functional architecture in the cat's visual cortex.** *J Physiol* 1962, **160**:106-154.
 3. Reid RC, Alonso JM: **The processing and encoding of information in the visual cortex.** *Curr Opin Neurobiol* 1996, **6**:475-480.
 4. Ferster D, Miller KD: **Neural mechanisms of orientation selectivity in the visual cortex.** *Annu Rev Neurosci* 2000, **23**:441-471.
 5. Reid RC, Alonso JM: **Specificity of monosynaptic connections from thalamus to visual cortex.** *Nature* 1995, **378**:281-284.
 6. Ferster D, Chung S, Wheat H: **Orientation selectivity of thalamic input to simple cells of cat visual cortex.** *Nature* 1996, **380**:249-252.
 7. Chung S, Ferster D: **Strength and orientation tuning of the thalamic input to simple cells revealed by electrically evoked cortical suppression.** *Neuron* 1998, **20**:1177-1189.
 8. Kara P, Pezaris JS, Yurgenson S, Reid RC: **The spatial receptive field of thalamic inputs to single cortical simple cells revealed by the interaction of visual and electrical stimulation.** *Proc Natl Acad Sci USA* 2002, **99**:16261-16266.
 9. Alonso JM, Martinez LM: **Functional connectivity between simple cells and complex cells in cat striate cortex.** *Nat Neurosci* 1998, **1**:395-403.
 10. Martinez LM, Wang Q, Reid RC, Pillai C, Alonso JM, Sommer FT, Hirsch JA: **Receptive field structure varies with layer in the primary visual cortex.** *Nat Neurosci* 2005, **8**:372-379.
 11. Mooser F, Bosking WH, Fitzpatrick D: **A morphological basis for orientation tuning in primary visual cortex.** *Nat Neurosci* 2004, **7**:872-879.
 12. LeVay S, Hubel DH, Wiesel TN: **The pattern of ocular dominance columns in macaque visual cortex revealed by a reduced silver stain.** *J Comp Neurol* 1975, **159**:559-576.
 13. Livingstone MS, Hubel DH: **Anatomy and physiology of a color system in the primate visual cortex.** *J Neurosci* 1984, **4**:309-356.
 14. Sincich LC, Horton JC: **Divided by cytochrome oxidase: a map of the projections from V1 to V2 in macaques.** *Science* 2002, **295**:1734-1737.
 15. Gilbert CD, Wiesel TN: **Intrinsic connectivity and receptive field properties in visual cortex.** *Vision Res* 1985, **25**:365-374.
 16. Lund JS: **Anatomical organization of macaque monkey striate visual cortex.** *Annu Rev Neurosci* 1988, **11**:253-288.
 17. Gilbert CD, Wiesel TN: **Columnar specificity of intrinsic horizontal and corticocortical connections in cat visual cortex.** *J Neurosci* 1989, **9**:2432-2442.
 18. Bosking WH, Zhang Y, Schofield B, Fitzpatrick D: **Orientation selectivity and the arrangement of horizontal connections in tree shrew striate cortex.** *J Neurosci* 1997, **17**:2112-2127.
 19. Braitenberg V, Schüz A: **Peters' rule and White's exceptions.** In *Cortex: Statistics and Geometry of Neuronal Connectivity*. Springer; 1998. pp. 99-102.
 20. Kalisman N, Silberberg G, Markram H: **The neocortical microcircuit as a tabula rasa.** *Proc Natl Acad Sci USA* 2005, **102**:880-885.
- Axons of layer 5 pyramidal neurons contacted the dendrites of neighboring layer 5 neurons without any bias. This supports that Peters' rule is correct for the geometrical contact between axons and dendrites. Functional connectivity as measured physiologically was correlated with the number of synaptic boutons at the contact sites.
21. Stepanyants A, Chklovskii DB: **Neurogeometry and potential synaptic connectivity.** *Trends Neurosci* 2005, **28**:387-394.
 22. Peters A: **Thalamic input to the cerebral cortex.** *Trends Neurosci* 1979, **2**:183-185.
 23. DeFelipe J, Elston GN, Fujita I, Fuster J, Harrison KH, Hof PR, Kawaguchi Y, Martin KA, Rockland KS, Thomson AM *et al.*: **Neocortical circuits: evolutionary aspects and specificity versus non-specificity of synaptic connections. Remarks, main conclusions and general comments and discussion.** *J Neurocytol* 2002, **31**:387-416.
 24. Shepherd GM, Stepanyants A, Bureau I, Chklovskii D, Svoboda K: **Geometric and functional organization of cortical circuits.** *Nat Neurosci* 2005, **8**:782-790.
- The authors compared geometrical and functional connectivities in rat barrel cortex. Geometrical connectivity was obtained from anatomical reconstructions of neurons and functional connectivity was obtained with photo-uncaging of glutamate. Functional connectivity could not be predicted solely from the overlap of dendrites and axonal arbors, but much of the variation could be explained when neurons' locations in layers and cortical maps were considered.
25. Mason A, Nicoll A, Stratford K: **Synaptic transmission between individual pyramidal neurons of the rat visual cortex in vitro.** *J Neurosci* 1991, **11**:72-84.
 26. Thomson AM, Deuchars J: **Synaptic interactions in neocortical local circuits: dual intracellular recordings in vitro.** *Cereb Cortex* 1997, **7**:510-522.
 27. Markram H, Lubke J, Frotscher M, Roth A, Sakmann B: **Physiology and anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat neocortex.** *J Physiol* 1997, **500**(Pt 2):409-440.
 28. Thomson AM, Bannister AP: **Interlaminar connections in the neocortex.** *Cereb Cortex* 2003, **13**:5-14.
 29. Holmgren C, Harkany T, Svennenfors B, Zilberter Y: **Pyramidal cell communication within local networks in layer 2/3 of rat neocortex.** *J Physiol* 2003, **551**:139-153.
 30. Song S, Sjöström PJ, Reigl M, Nelson S, Chklovskii DB: **Highly nonrandom features of synaptic connectivity in local cortical circuits.** *PLoS Biol* 2005, **3**:e68.
 31. Yoshimura Y, Dantzker JL, Callaway EM: **Excitatory cortical neurons form fine-scale functional networks.** *Nature* 2005, **433**:868-873.
- The authors showed the existence of subnetworks embedded in larger functional architecture in rat visual cortex. Neighboring layer 2/3 pyramidal neurons were functionally connected with each other only when they received common inputs from layer 4.
32. Yoshimura Y, Callaway EM: **Fine-scale specificity of cortical networks depends on inhibitory cell type and connectivity.** *Nat Neurosci* 2005, **8**:1552-1559.
 33. Blasdel GG, Salama G: **Voltage-sensitive dyes reveal a modular organization in monkey striate cortex.** *Nature* 1986, **321**:579-585.
 34. Bonhoeffer T, Grinvald A: **Iso-orientation domains in cat visual cortex are arranged in pinwheel-like patterns.** *Nature* 1991, **353**:429-431.
 35. Polimeni JR, Granquist-Fraser D, Wood RJ, Schwartz EL: **Physical limits to spatial resolution of optical recording: clarifying the spatial structure of cortical hypercolumns.** *Proc Natl Acad Sci USA* 2005, **102**:4158-4163.

36. Stosiek C, Garaschuk O, Holthoff K, Konnerth A: **In vivo two-photon calcium imaging of neuronal networks.** *Proc Natl Acad Sci USA* 2003, **100**:7319-7324.
37. Svoboda K, Denk W, Kleinfeld D, Tank DW: **In vivo dendritic calcium dynamics in neocortical pyramidal neurons.** *Nature* 1997, **385**:161-165.
38. Ohki K, Chung S, Ch'ng YH, Kara P, Reid RC: **Functional imaging with cellular resolution reveals precise micro-architecture in visual cortex.** *Nature* 2005, **433**:597-603.
In vivo two-photon calcium imaging was used to map the orientation/direction selectivity of thousands of neurons at single-cell resolution in rat and cat visual cortices. In the rat visual cortex, neurons were not organized for orientation. In the cat visual cortex, neurons were arranged very precisely according to their preferred orientation and direction, and the functional border between direction domains was very precise.
39. Powell TP, Mountcastle VB: **Some aspects of the functional organization of the cortex of the postcentral gyrus of the monkey: a correlation of findings obtained in a single unit analysis with cytoarchitecture.** *Bull Johns Hopkins Hosp* 1959, **105**:133-162.
40. Favorov OV, Diamond ME: **Demonstration of discrete place-defined columns – segregates – in the cat SI.** *J Comp Neurol* 1990, **298**:97-112.
41. Hubel DH, Wiesel TN: **Sequence regularity and geometry of orientation columns in the monkey striate cortex.** *J Comp Neurol* 1974, **158**:267-293.
42. Ohki K, Chung S, Kara P, Hübener M, Bonhoeffer T, Reid RC: **Highly ordered arrangement of single neurons in orientation pinwheels.** *Nature* 2006, **442**:925-928.
43. Shmuel A, Grinvald A: **Functional organization for direction of motion and its relationship to orientation maps in cat area 18.** *J Neurosci* 1996, **16**:6945-6964.
44. Maldonado PE, Godecke I, Gray CM, Bonhoeffer T: **Orientation selectivity in pinwheel centers in cat striate cortex.** *Science* 1997, **276**:1551-1555.
45. Schummers J, Marino J, Sur M: **Synaptic integration by V1 neurons depends on location within the orientation map.** *Neuron* 2002, **36**:969-978.
46. Peters A, Sethares C: **Organization of pyramidal neurons in area 17 of monkey visual cortex.** *J Comp Neurol* 1991, **306**:1-23.
47. Peters A, Yilmaz E: **Neuronal organization in area 17 of cat visual cortex.** *Cereb Cortex* 1993, **3**:49-68.
48. Marino J, Schummers J, Lyon DC, Schwabe L, Beck O, Wiesing P, Obermayer K, Sur M: **Invariant computations in local cortical networks with balanced excitation and inhibition.** *Nat Neurosci* 2005, **8**:194-201.
49. Chapman B, Stryker MP, Bonhoeffer T: **Development of orientation preference maps in ferret primary visual cortex.** *J Neurosci* 1996, **16**:6443-6453.
50. Li Y, Fitzpatrick D, White LE: **The development of direction selectivity in ferret visual cortex requires early visual experience.** *Nat Neurosci* 2006, **9**:676-681.
51. Hubel DH, Wiesel TN: **Shape and arrangement of columns in cat's striate cortex.** *J Physiol* 1963, **165**:559-568.
52. Van Hooser SD, Heimel JA, Chung S, Nelson SB, Toth LJ:
 • **Orientation selectivity without orientation maps in visual cortex of a highly visual mammal.** *J Neurosci* 2005, **25**:19-28.
 The gray squirrel has a relatively large visual cortex and good visual acuity compared with smaller rodents. However, the authors showed that squirrel's visual cortex was not organized for orientation, though individual neurons were highly orientation selective.
53. Adams DL, Horton JC: **Monocular cells without ocular dominance columns.** *J Neurophysiol* 2006, **96**:2253-2264.
54. Horton JC, Adams DL: **The cortical column: a structure without a function.** *Philos Trans R Soc Lond B Biol Sci* 2005, **360**:837-862.
 A historical perspective on the significance of functional columns. The authors suggest that there is no evident relationship between existences of functional columns and visual function.
55. Dräger UC: **Receptive fields of single cells and topography in mouse visual cortex.** *J Comp Neurol* 1975, **160**:269-290.
56. Mangini NJ, Pearlman AL: **Laminar distribution of receptive field properties in the primary visual cortex of the mouse.** *J Comp Neurol* 1980, **193**:203-222.
57. Metin C, Godement P, Imbert M: **The primary visual cortex in the mouse: receptive field properties and functional organization.** *Exp Brain Res* 1988, **69**:594-612.
58. Wiesenfeld Z, Kornel EE: **Receptive fields of single cells in the visual cortex of the hooded rat.** *Brain Res* 1975, **94**:401-412.
59. Parnavelas JG, Burne RA, Lin CS: **Receptive field properties of neurons in the visual cortex of the rat.** *Neurosci Lett* 1981, **27**:291-296.
60. Girman SV, Sauve Y, Lund RD: **Receptive field properties of single neurons in rat primary visual cortex.** *J Neurophysiol* 1999, **82**:301-311.
61. Wang KH, Majewska A, Schummers J, Farley B, Hu C, Sur M, Tonegawa S: **In vivo two-photon imaging reveals a role of arc in enhancing orientation specificity in visual cortex.** *Cell* 2006, **126**:389-402.
62. Sohya K, Kameyama K, Yanagawa Y, Obata K, Tsumoto T:
 • **GABAergic neurons are less selective to stimulus orientation than excitatory neurons in layer II/III of visual cortex, as revealed by in vivo functional Ca2+ imaging in transgenic mice.** *J Neurosci* 2007, **27**:2145-2149.
 Combining *in vivo* two-photon calcium imaging with transgenic mice expressing GFP in GABAergic neurons, the authors showed that GABAergic neurons were less tuned for orientation than pyramidal neurons in visual cortex.
63. Van Hooser SD, Heimel JA, Chung S, Nelson SB: **Lack of patchy horizontal connectivity in primary visual cortex of a mammal without orientation maps.** *J Neurosci* 2006, **26**:7680-7692.
64. Koulakov AA, Chklovskii DB: **Orientation preference patterns in mammalian visual cortex: a wire length minimization approach.** *Neuron* 2001, **29**:519-527.
65. Denk W, Horstmann H: **Serial block-face scanning electron microscopy to reconstruct three-dimensional tissue nanostructure.** *PLoS Biol* 2004, **2**:e329.
66. DeFalco J, Tomishima M, Liu H, Zhao C, Cai X, Marth JD, Enquist L, Friedman JM: **Virus-assisted mapping of neural inputs to a feeding center in the hypothalamus.** *Science* 2001, **291**:2608-2613.
67. Kelly RM, Strick PL: **Cerebellar loops with motor cortex and prefrontal cortex of a nonhuman primate.** *J Neurosci* 2003, **23**:8432-8444.
68. Wickersham IR, Finke S, Conzelmann KK, Callaway EM: **Retrograde neuronal tracing with a deletion-mutant rabies virus.** *Nat Methods* 2007, **4**:47-49.
69. Wickersham IR, Lyon DC, Barnard RJ, Mori T, Finke S,
 • • Conzelmann KK, Young JA, Callaway EM: **Monosynaptic restriction of transsynaptic tracing from single, genetically targeted neurons.** *Neuron* 2007, **53**:639-647.
 The authors developed a modified rabies virus, which crosses only one synaptic step retrogradely. Using this virus, they identified all the neurons providing monosynaptic inputs to a single neuron. This technique will help reveal the detailed connectivity of the cortical circuits.
70. Oliva AA Jr, Jiang M, Lam T, Smith KL, Swann JW: **Novel hippocampal interneuronal subtypes identified using transgenic mice that express green fluorescent protein in GABAergic interneurons.** *J Neurosci* 2000, **20**:3354-3368.
71. Meyer AH, Katona I, Blatow M, Rozov A, Monyer H: **In vivo labeling of parvalbumin-positive interneurons and analysis of electrical coupling in identified neurons.** *J Neurosci* 2002, **22**:7055-7064.
72. Chattopadhyaya B, Di Cristo G, Higashiyama H, Knott GW, Kuhlman SJ, Welker E, Huang ZJ: **Experience and activity-dependent maturation of perisomatic GABAergic innervation in primary visual cortex during a postnatal critical period.** *J Neurosci* 2004, **24**:9598-9611.

73. Lopez-Bendito G, Sturgess K, Erdelyi F, Szabo G, Molnar Z, Paulsen O: **Preferential origin and layer destination of GAD65-GFP cortical interneurons.** *Cereb Cortex* 2004, **14**:1122-1133.
74. Dumitriu D, Cossart R, Huang J, Yuste R: **Correlation between axonal morphologies and synaptic input kinetics of interneurons from mouse visual cortex.** *Cereb Cortex* 2007, **17**:81-91.
75. Lechner HA, Lein ES, Callaway EM: **A genetic method for selective and quickly reversible silencing of mammalian neurons.** *J Neurosci* 2002, **22**:5287-5290.
76. Karpova AY, Tervo DG, Gray NW, Svoboda K: **Rapid and reversible chemical inactivation of synaptic transmission in genetically targeted neurons.** *Neuron* 2005, **48**:727-735.
77. Tan EM, Yamaguchi Y, Horwitz GD, Gosgnach S, Lein ES, Goulding M, Albright TD, Callaway EM: **Selective and quickly reversible inactivation of mammalian neurons *in vivo* using the *Drosophila* allatostatin receptor.** *Neuron* 2006, **51**:157-170.
- Genetic silencing of neurons with allatostatin receptor was tested *in vivo* in thalamic and cortical neurons of rat, ferret, and monkey. Application of its ligand dramatically decreased neural activity, and the inactivation was reversible within minutes.
78. Gosgnach S, Lanuza GM, Butt SJ, Saueressig H, Zhang Y, Velasquez T, Riethmacher D, Callaway EM, Kiehn O, Goulding M: **V1 spinal neurons regulate the speed of vertebrate locomotor outputs.** *Nature* 2006, **440**:215-219.
79. Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K: **•• Millisecond-timescale, genetically targeted optical control of neural activity.** *Nat Neurosci* 2005, **8**:1263-1268.
- A light-gated cation channel, channelrhodopsin-2, was introduced into neurons, thus allowing individual action potentials to be evoked by light with millisecond resolution. This study opened up the possibility of controlling activity in subclasses of neurons when channelrhodopsin is expressed under a specific promoter. Subsequent studies have proven the usefulness of this tool for studying cortical circuitry.
80. Han X, Boyden ES: **Multiple-color optical activation, silencing, and desynchronization of neural activity, with single-spike temporal resolution.** *PLoS ONE* 2007, **2**:e299.
- A chloride channel, halorhodopsin, that is excited by yellow light was used to inhibit firing of neurons. When combined with channelrhodopsin-2, excited by blue light, individual action potentials could be added or deleted from spike trains of single neurons.
81. Zhang F, Wang LP, Brauner M, Liewald JF, Kay K, Watzke N, Wood PG, Bamberg E, Nagel G, Gottschalk A *et al.*: **Multimodal fast optical interrogation of neural circuitry.** *Nature* 2007, **446**:633-639.
- A light-gated cation channel (channelrhodopsin-2) and chloride channel (NpHR, halorhodopsin) were introduced into neurons so that they could be excited and inhibited by light. In cortical slices, the effects of light-gated channels were imaged with calcium indicators. In *C. elegans*, the channels were used to control behavior.